

RSC Communicable and Respiratory Disease Report for England

Week Number / Year 21 / 2025	Population 16,580,080
Dates 19/05/2025 - 25/05/2025	No. Practices 1,567

Notes

All rates in this report are given per 100,000 population presenting in the week of the report. A rolling 5-year average rate is also provided as a historical comparison. Rates are provided for four regions (North, South, Midlands and East, and London). For acute respiratory infections, a breakdown by age group is also provided.

Rates are presented on a weekly basis, using ISO week numbers.

Please see page 20 for further explanatory notes on the data.

Comments

Overall rates of influenza-like illness (ILI) are stable in all regions and are around the seasonal average for this time of year (pages 3 to 5). ILI rates are now below the medium threshold across all age bands: see Table (E), page 5.

Rates of acute respiratory infections (ARI) are stable across all regions, remaining at or below the seasonal average, page 7.

Overall rates of COVID-19 are stable and remain low, page 6.

This report includes a respiratory virology update: see Graph (C), page 4.

Other comments:

• Rates of measles in London continue to increase (page 15) despite the national rate of measles remaining at the national average for this time of year and the overall number of cases of measles falling slightly from week 20 (Page 19).

** Rates of scabies (page 16) remain above the seasonal average.

Seasonal Focus

In the "Change since last week" column, a change in rate of 5% to 10% is marked with a single arrow (\land or \checkmark), while a change of more than 10% is marked with a double arrow (\land or \checkmark). A flat line (-) indicates the rate was stable, changing less than 5%.

Region Breakdown

	Acute respiratory infections (ARI)			Influe	nza-like	illness (ILI)	Exacerbations of chronic lung disease (ECLD)			
	This week	Last week	Change since last week	This week	Last week	Change since last week	This week	Last week	Change since last week	
London	153.2	151.20	- 1.9	2.2	2.30	✓ -0.1	8.6	9.38	∨ -0.8	
Midlands And East	182.9	181.20	- 1.7	1.2	0.9	☆ 0.2	13.5	13.30	- 0.2	
North	226.8	230.0	- -3.2	1.8	1.8	- 0.0	19.4	20.3	- -1.0	
South	163.0🛛	172.50	∽-9.5	1.5	1.8	♥ -0.3	11.8	12.4	- -0.6	
National	181.3	184.5	- -3.2	1.6	1.7	- -0.1	13.4	14.0	- -0.6	
	Lower respiratory tract infections (LRTI)		Upper respiratory tract infections (URTI)				COVID	-19		
	in	fections	(LRTI)	in	fections	(URTI)				
	in This week	fections Last week	(LRTI) Change since last week	in This week	fections Last week	(URTI) Change since last week	This week	Last week	Change since last week	
London	in This week 37.2⊠	fections Last week 37.2	(LRTI) Change since last week - 0.0	in This week	fections Last week 108.5	(URTI) Change since last week - 1.8	This week	Last week	Change since last week	
London Midlands And East	in This week 37.20 58.10	fections Last week 37.2 60.3	(LRTI) Change since last week - 0.0 2.2	in This week 110.30 116.90	fections Last week 108.5 113.9	(URTI) Change since last week - 1.8 - 3.0	This week 0.70 0.50	Last week 1.0⊠ 0.6	Change since last week	
London Midlands And East North	in This week 37.20 58.10 79.00	fections Last week 37.21 60.31 80.11	(LRTI) Change since last week - 0.0 2.2 1.1	in This week 110.30 116.90 138.40	fections Last week 108.5% 113.9% 140.7%	(URTI) Change since last week - 1.8 - 3.0 2.3	This week 0.71/2 0.55/2 0.8	Last week 1.01 0.6 0.91	Change since last week	
London Midlands And East North South	in This week 37.20 58.10 79.00 54.6	fections Last week 37.20 60.30 80.10 57.40	(LRTI) Change since last week - 0.0 2.2 1.1 2.8	in This week 110.3 116.9 138.4 104.1	fections Last week 108.5% 113.9% 140.7% 109.6	(URTI) Change since last week - 1.8 - 3.0 2.3 5.5	This week 0.7№ 0.5№ 0.8 0.8	Last week 1.0⊠ 0.6 0.9⊠ 0.8	Change since last week	

Age Group Breakdown

	Acute respiratory infections (ARI)		Influ	enza-like	illness (ILI)	Exacerbations of chronic lung disease (ECLD)				
	This week	Last week	Change since last week	This week	Last week	Change since last week	This week	Last week	Change since last week	
<1yr	908.6	914.0	- -5.4	1.5	2.3	癸 -0.8	0.0	0.0⊠	- 0.0	
1-4yrs	628.4	655.1🛛	- -26.6	1.6	2.3	癸 -0.7	0.9⊠	2.8	≽ -1.8	
5-14yrs	219.3	221.4	- -2.2	1.4	0.7⊠	♠ 0.6	6.3⊠	6.2	- 0.1	
15-64yrs	130.4	132.8	- -2.5	1.8	2.0	癸-0.2	9.8	10.0⊠	- -0.2	
65+yrs	216.0	218.7	- -2.7	1.30	1.1⊠	♠ 0.2	34.8	36.6	- -1.8	
All ages	181.3	184.5	- -3.2	1.6	1.7	- -0.1	13.4	14.0	- -0.6	
Lower respiratory tract infections			Upper respiratory tract infections				COVID-19			

(LRTI) (URTI) This Last Change since This Last Change since This Last Change since week week last week week week last week week week last week 811.8 <1yr 154.20 178.0 **४** −23.8 796.8 - 14.9 4.5 5.30 ₿-0.8 1-4yrs 87.2 94.7 ✓ -7.5 574.1 591.18 --17.0 0.3 0.30 -0.0 5-14yrs 25.10 25.6 **-** -0.5 193.10 194.6 - -1.6 0.10 ¥-0.1 0.2 ♥-0.1 15-64yrs 40.6 41.4 **-** -0.8 84.5 85.8 - -1.3 0.6 0.70 65+yrs 133.10 136.9 - -3.8 55.8 56.3 --0.6 1.50 1.70 ₩-0.2 All ages 57.9 59.7 116.8 118.1 - -1.3 0.7 0.8 **≫** -0.1

2024/25 Focus

(A) Influenza-like Illness: national incidence rate by region

The horizontal lines in the following graph are thresholds derived from the Moving Epidemic Method (MEM) model. See p20 for more information.



(B) RCGP/UKHSA influenza virology swab surveillance





(C) RCGP/UKHSA RSV, influenza and SARS-CoV-2 virology swab surveilance (by strain)

(D) Influenza-like Illness: national incidence rate by age band



(E) Influenza-like Illness: national incidence rate by age band

This table shows the level of intensity of ILI by age band. MEM thresholds have been calculated separately for each age band - thresholds are shown in the second table. Refer to page 19 for more information.

	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	2	1	2	3
1-4yrs	3.0	2.1	2.2	2.6	4.2	4.3	4.0	1.9	4.7	6.6	11.0	11.1	18.9	34.2	24.	7 2	27.5	22.0	24.0
5-14yrs	1.2	1.8	1.5	1.5	1.7	2.0	1.3	1.9	2.9	3.4	4.4	6.8	9.2	14.4	8.	1	6.1	8.2	10.1
15-64yrs	2.5	2.8	3.6	4.0	4.4	4.1	4.1	4.3	4.1	5.0	6.2	9.5	13.1	18.1	13.	2 2	20.2	24.8	18.0
65+yrs	1.8	1.7	2.9	3.3	3.7	3.5	3.3	3.4	2.8	2.7	5.6	5.2	8.5	15.5	17.	1 2	29.3	26.4	16.1
All ages	2.3	2.5	3.2	3.6	3.9	3.8	3.6	3.8	3.7	4.5	6.0	8.5	12.1	18.0	13.	9 2	20.6	23.1	17.0
	4		5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21
1-4yrs	21.3	22	.1 1	8.4	13.5	11.9	6.9	8.8	7.0	4.8	5.8	5.4	4.2	1.7	2.0	2.0	1.1	2.3	1.6
5-14yrs	12.8	13	.8 1	1.3	9.3	6.7	4.9	5.0	4.6	5.2	4.6	2.9	1.7	0.8	0.9	0.8	0.7	0.7	1.4
15-64yrs	15.8	14	.5 1	3.5	12.2	11.5	10.6	8.8	7.6	7.0	6.2	5.2	4.0	2.8	2.5	2.2	1.7	2.0	1.8
65+yrs	14.3	10	.0	8.8	8.9	6.3	6.2	5.5	3.9	4.0	3.8	4.4	4.9	2.2	1.9	1.8	1.3		1.3
All ages	15.4	13	.9 1	2.6	11.3	10.1	9.1	7.8	6.6	6.2	5.6	4.8	4.0	2.4	2.2	2.0	1.5	1.7	1.6
	Polow	Thr	achal	d	Throck	and to	Mod	ium	Mod	diuma	to High		iah to	Vorv	High	٨		Vory L	liah

	Below Threshold	Threshold to Medium	Medium to High	High to Very High	Above Very High
1-4yrs	<7.9	7.9 to 12.6	12.6 to 26.2	26.2 to 36.1	36.1+
5-14yrs	<5.4	5.4 to 10.7	10.7 to 26.6	26.6 to 39.9	39.9+
15-64yrs	<9.8	9.8 to 17.9	17.9 to 43.0	43.0 to 63.4	63.4+
65+yrs	<9.3	9.3 to 15.0	15.0 to 38.8	38.8 to 59.0	59.0+
All Ages	<8.54	8.54 to 16.27	16.27 to 38.66	38.66 to 56.68	56.68+

(F) Acute Bronchitis and Bronchiolitis: national incidence rate by age band

Children under 1 year old are omitted from the following graph.



Weekly incidence rates of influenza-like illness, and acute bronchitis and bronchiolitis (per 100,000)

	Influenza-like illness (ILI)	ARI-Bronchitis and Bronchiolitis		Influenza-like illness (ILI)	ARI-Bronchitis and Bronchiolitis
<1vr	1.5	81.90	London	2.2	1.5
1-4yrs	1.6	8.6	Midlands And East	1.2	2.0
5-14yrs	1.4	0.5	North	1.8	2.4
15-24yrs	1.2	0.3	South	1.5	2.0
25-44yrs	2.1	0.8	National	1.6	2.0
45-64yrs	1.6	1.10			
65-74yrs	1.0	2.1			
75-84yrs	1.3🛙	2.4⊠			
85+yrs	2.01	2.2			
All ages	1.6	2.0			

1.50 2.0 2.40 2.0 2.0



(G) COVID-19: national incidence rate by region

(H) COVID-19: national incidence rate by age band



1. Respiratory Infections



(I) Acute Respiratory Infections (ARI): national incidence rate by region

(J) Acute Respiratory Infections (ARI): national incidence rate by age band



Respiratory Infections - by region



London South





Exacerbations of Chronic Lung Disease (ECLD) Weekly incidence (per 100,000 all ages) by region for 2024/25 compared with 5 year average



Lower Respiratory Tract Infections (LRTI) Weekly incidence (per 100,000 all ages) by region for 2024/25 compared with 5 year average



Respiratory Infections - by age band



Influenza-like illness (ILI) Weekly incidence (per 100,000 all ages) by age band for 2024/25 compared with 5 year average



Exacerbations of Chronic Lung Disease (ECLD) Weekly incidence (per 100,000 all ages) by age band for 2024/25 compared with 5 year average



Lower Respiratory Tract Infections (LRTI) Weekly incidence (per 100,000 all ages) by age band for 2024/25 compared with 5 year average



Respiratory Infections - by region



London South

Upper Respiratory Tract Infections (URTI) Weekly incidence (per 100,000 all ages) by region for 2024/25 compared with 5 year average



ECLD - Asthma Exacerbations Weekly incidence (per 100,000 all ages) by region for 2024/25 compared with 5 year average



ECLD - COPD Exacerbations Weekly incidence (per 100,000 all ages) by region for 2024/25 compared with 5 year average



Respiratory Infections - by age band



Upper Respiratory Tract Infections (URTI) Weekly incidence (per 100,000 all ages) by age band for 2024/25 compared with 5 year average



ECLD - Asthma Exacerbations Weekly incidence (per 100,000 all ages) by age band for 2024/25 compared with 5 year average



ECLD - COPD Exacerbations Weekly incidence (per 100,000 all ages) by age band for 2024/25 compared with 5 year average



All ages

5 Year Avg

Respiratory Infections - by region

Respiratory Infections - by age band

<1yr

1-4yrs



London South





LRTI - Acute Bronchitis Weekly incidence (per 100,000 all ages) by region for 2024/25 compared with 5 year average



LRTI - Bronchiolitis Weekly incidence (per 100,000 all ages) by region for 2024/25 compared with 5 year average



LRTI - Pneumonia

5-14yrs

15-64yrs

65+yrs

Weekly incidence (per 100,000 all ages) by age band for 2024/25 compared with 5 year average



LRTI - Acute Bronchitis Weekly incidence (per 100,000 all ages) by age band for 2024/25 compared with 5 year average



LRTI - Bronchiolitis Weekly incidence (per 100,000 all ages) by age band for 2024/25 compared with 5 year average



Respiratory Infections - by region

National 5 Year Avg North Midlands And East

London South





URTI - Tonsillitis/Pharyngitis Weekly incidence (per 100,000 all ages) by region for 2024/25 compared with 5 year average



URTI - Otitis Media Weekly incidence (per 100,000 all ages) by region for 2024/25 compared with 5 year average



All ages <1yr 5-14yrs 65+yrs 5 Year Avg 1-4yrs 15-64yrs

Respiratory Infections - by age band

URTI - Croup Weekly incidence (per 100,000 all ages) by age band for 2024/25 compared with 5 year average



URTI - Tonsillitis/Pharyngitis Weekly incidence (per 100,000 all ages) by age band for 2024/25 compared with 5 year average



URTI - Otitis Media Weekly incidence (per 100,000 all ages) by age band for 2024/25 compared with 5 year average



Respiratory Infections - by region



🗾 London 📒 South





URTI - Laryngitis Weekly incidence (per 100,000 all ages) by region for 2024/25 compared with 5 year average



Respiratory Infections - by age band



URTI - Sinusitis Weekly incidence (per 100,000 all ages) by age band for 2024/25 compared with 5 year average



URTI - Laryngitis Weekly incidence (per 100,000 all ages) by age band for 2024/25 compared with 5 year average





0.25

0.00

40 44 48 52 4 8 12 16 20 24 28 32 36

1

0+ 40

44 48 52 4 8 12 16 20 24 28 32 36







Page 16 of 22

0+

12 16 20 24 28 32 36

48 52 4 8 12 16 20 24 28 32 36



6. Disorders Affecting the Nervous System

Disorders of Peripheral Nervous System (ICD10: G50-G64,G70-G72), Weekly incidence (per 100,000 all ages) by region for 2024/25 compared with 5 year average



Symptoms of Nervous & Musculoskeletal Systems (ICD10: R25-R29), Weekly incidence (per 100,000 all ages) by region for 2024/25 compared with 5 year average



Meningitis/Encephalitis (ICD10: A170-A171,A390,A38-A85,A87,G00-G05), Weekly incidence (per 100,000 all ages) by region for 2024/25 compared with 5 year average







8. Other Disorders

Strep Sore Throat, Scarlatina and Peritonsillar Abscess (ICD10: A38,J020,J36), Weekly incidence (per 100,000 all ages) by region for 2024/25 compared with 5 year average







9. Tabular Summary by Disease

	Week 18	Week 19	Week 20	Week 21
Dates	28/04/2025 - 04/05/2025	05/05/2025 - 11/05/2025	12/05/2025 - 18/05/2025	19/05/2025 - 25/05/2025
Population	16,887,804	17,209,933	16,994,963	16,580,080
Practice Count	1,580	1,608	1,596	1,567

	Wee	k 18	Wee	ek 19	Wee	ek 20	Wee	ek 21
Disease	Rate	Count	Rate	Count	Rate	Count	Rate	Count
Acute Bronchitis	1.4	228	1.0	171	1.1	183	1.0	174
Acute Respiratory Infections (ARI)	179.9	30,379	159.8	27,502	184.5	31,360	181.3	30,066
Allergic Rhinitis	22.7	3,830	15.8	2,715	22.2	3,779	18.7	3,107
Asthma	10.2	1,721	7.8	1,334	10.2	1,740	10.1	1,672
Bronchiolitis	0.9	146	0.9	163	1.1	182	0.9	155
Bullous Dermatoses	0.2	27	0.2	38	0.1	23	0.2	35
COVID-19	0.7	124	0.5	90	0.8	139	0.7	117
Chickenpox	4.0	672	3.5	602	3.7	631	3.7	617
Conjunctival Disorders	14.8	2,492	12.0	2,065	15.2	2,591	14.3	2,371
Croup	1.5	247	1.4	236	1.9	319	1.6	265
ECLD - COPD exacerbations	6.0	1,010	5.1	883	5.6	951	5.4	893
ECLD - asthma exacerbations	8.5	1,433	7.2	1,243	8.5	1,445	8.1	1,343
Exacerbations of chronic lung disease (ECLD)	14.4	2,433	12.2	2,099	14.0	2,374	13.4	2,223
Herpes Simplex	2.7	458	2.8	475	3.6	611	2.9	481
Herpes Zoster	5.0	852	4.2	729	4.9	827	4.8	797
Impetigo	2.7	453	2.7	460	2.9	496	2.7	440
Infected Insect Bites	3.9	652	2.9	505	5.7	962	4.4	730
Infectious Intestinal Diseases	7.3	1,231	6.7	1,161	8.8	1,504	8.4	1,385
Infectious Mononucleosis	0.3	56	0.3	52	0.3	58	0.3	49
Influenza-like Illness (ILI)	2.0	331	1.5	260	1.7	289	1.6	270
Laryngitis	0.8	131	0.8	130	0.8	133	0.7	120
Lower respiratory tract infections (LRTI)	61.9	10,446	54.3	9,341	59.7	10,144	57.9	9,596
Measles	0.0	8	0.1	9	0.1	11	0.1	9
Meningitis and Encephalitis	0.1	15	0.1	22	0.2	26	0.1	18
Mumps	0.1	14	0.0	5	0.0	6	0.0	7
Non-infective Enteritis and Colitis	2.3	393	2.0	349	2.8	468	2.6	436
Peripheral Nervous Disease	18.4	3,108	15.7	2,705	20.4	3,463	19.1	3,167
Pneumonia	2.7	458	2.9	503	3.2	545	2.6	423
Rubella	0.0	0	0.0	0	0.0	1	0.0	1
Scabies	3.0	502	2.4	410	2.9	488	2.7	445
Sinusitis	15.7	2,647	13.6	2,335	14.8	2,515	14.8	2,459
Skin and Subcutaneous Tissue Infections	84.4	14,261	68.6	11,812	87.9	14,939	83.0	13,767
Strep Throat and Peritonsillar Abscess	1.0	169	1.2	200	1.1	192	1.2	191
Symptoms involving Skin and Integument Tissues	134.8	22,772	107.3	18,461	137.1	23,304	130.4	21,616
Symptoms involving musculoskeletal	14.6	2,460	13.1	2,253	16.0	2,726	14.9	2,478
Tonsillitis and Pharyngitis	27.0	4,553	23.6	4,066	28.2	4,800	27.8	4,612
Upper respiratory tract infections (URTI)	111.7	18,860	100.3	17,253	118.1	20,065	116.8	19,362
Urinary Tract Infections	22.6	3,809	23.1	3,976	25.3	4,307	23.2	3,846
Viral Hepatitis	0.2	41	0.2	31	0.3	51	0.2	34
Whooping Cough	0.0	8	0.0	5	0.0	7	0.0	6

Further Information

Focus on winter respiratory infections and infections with epidemic or pandemic infection

A key role of the RSC is to monitor conditions that cause winter pressures on the NHS, as well as provide early warnings of outbreaks, epidemics, and pandemics. The RSC has been collecting data on infections since 1957, conducting sentinel surveillance since 1967 (with virology added in 1993), and serosurveillance from 2000.

Pages 2-6 of this report focus on influenza-like illness (ILI), virology data, and acute respiratory infections (ARI). ILI is the name given to clinically identified flu cases, around half of which will be due to the influenza virus (the other half will be due to other viruses).

Measuring the level of circulating influenza

The level of influenza-like illness (ILI) is reported using intensity thresholds (Graph A, page 2 and Table E, page 4). These are calculated using the Moving Epidemic Method (MEM). MEM works by identifying seasonal epidemic peaks and then calculating a baseline threshold and intensity levels based on pre- and post-epidemic rates. This provides a better measure of severity of ILI than simply comparing it to the five-year average rate.

The MEM intensity levels for ILI are defined as follows:

Threshold to Medium Below 40% percentile

Medium to High	From 40% to below 90% percentile
High to Very High	From 90% to below 97.5% percentile
Above Very High	At or above 97.5% percentile

The MEM methodology is used by the UK Health Security Agency (UKHSA) and by the European Centre for Disease Prevention and Control (ECDC) to standardise reporting of influenza activity.

More information about MEM can be found at:

https://www.ecdc.europa.eu/en/news-events/acute-respiratory-infections-eueea-epidemiological-update-and-current-public-health

Rate of monitored conditions

Our monitored conditions are reported as the number of new cases each week per 100,000 population. We refer to this as the 'weekly incidence'. All conditions are shown with males and females combined.

The report's population, also called the denominator, is the registered population of RSC practices who share anonymised data for this report. The denominator varies weekly as patients register and deregister; additionally, a practice's data may not be included if there is an issue with data extraction.

Five-year averages

In addition to weekly incidence rates, we plot a five-year average for most conditions. Previously a ten-year average was used, but this window was shortened to reflect faster changes in seasonal variations and therefore enable a more meaningful comparison to relevant historic trends. COVID-19 pandemic years are excluded from this calculation for some conditions.

Regional rates of monitored conditions

In addition to a national rate, we present regional rates for all monitored conditions for four regions of England. The four RSC regions are aggregated NHS regions:

North	NHS North East and Yorkshire, and North West regions
Midlands and East	NHS East of England and Midlands regions
South	NHS South East and South West regions
London	NHS London region

Reporting of acute respiratory infections (ARI) by age band

In addition to regional rates, we report rates by age band for ARI. We display five age bands: those aged under 1 year, 1-4 years, 5-14 years, 15-64 years, and those aged 65 years and over. We subdivide ARI into four categories:

- influenza-like illness (ILI);
- exacerbations of chronic lung disease (ECLD), mainly asthma and chronic obstructive pulmonary disease (COPD);
- lower respiratory tract infections (LRTI), including bronchitis and pneumonia;
- upper respiratory tract infections (URTI), including tonsilitis and sinusitis.

More information about our classification of ARI can be found at:

https://www.eurosurveillance.org/content/10.2807/1560-7917.ES.2024.29.35.2300682

About the RCGP Research and Surveillance Centre (RSC)

What we do

Established in 1957, the Oxford-Royal College of General Practitioners (RCGP) Research and Surveillance Centre (RSC) is an internationally renowned source of information, analysis and interpretation concerning the onset, patterns, prevalence and trends over time of morbidity in primary care. The RSC provides weekly reports about health and disease: the Weekly Returns Service (WRS). The WRS has been produced since 1967, in collaboration with the UK Health Security Agency (UKHSA) and its predecessor bodies. The University of Oxford currently provides the WRS on behalf of RCGP and UKHSA.

The RSC is active in research and surveillance. In addition to the WRS, the RSC contributes data to UKHSA's Syndromic Surveillance system, and supports vaccine effectiveness studies. The role of general practice members of the RSC is set out in an annual commissioning letter.

Further information about the RSC can be found on our website:

www.rcgp.org.uk/representing-you/research-at-rcgp/research-surveillance-centre

Our data extraction process and governance

Data are extracted on behalf of the RSC from practice computerised medical record systems, twice a week by Magentus Data Management, or daily by EMIS-X Analytics (EXA).

Data are pseudonymised as close to source as possible. Data are held on secure servers at the Nuffield Department of Primary Care Health Sciences (NDPCHS) at the University of Oxford. Our systems meet the requirements of the General Data Protection Regulation (GDPR). Further information about the NHS England approval of the RSC's data security can be found at:

https://www.dsptoolkit.nhs.uk/OrganisationSearch/EE133863-MSD-NDPCHS

What the data is used for

The WRS monitors the number of patients consulting with new episodes of illness classified by diagnosis in England and provides weekly incidence rates per 100,000 population for these new episodes of illness. It is the key primary care element of the national disease monitoring systems run by the UK Health Security Agency.

In addition to the WRS, the data are used for other research studies. Any other uses of the data for research follow ethical approval or agreement from NIHR proportionate review, and where relevant Health Research Authority Confidential Advisory Group advice that further approval is not needed.

Get in touch

For further information about the work of the RSC, or if you would like to be included on our email notification list, please contact:

Director: Professor Simon de Lusignan (Simon.DeLusignanPA@phc.ox.ac.uk)

RCGP Research and Surveillance Centre Policy, Research and Campaigns Royal College of General Practitioners 30 Euston Square London, NW1 2FB Tel: 020 3188 7400



Nuffield Department of Primary Care Health Sciences Gibson Building Radcliffe Observatory Quarter Woodstock Road Oxford, OX2 6GG Tel: 01865 617855

